

Linear Systems and Signals

Steps, ramps, and rects

Anand D. Sarwate

Department of Electrical and Computer Engineering
Rutgers, The State University of New Jersey

2020




Learning objectives

The learning objectives for this section are:

- understand the definitions of the step, rect, and ramp functions
- use the unit step function to make signals right or left-sided
- use the rectangle function and its shifts/dilations to window signals



The unit step function



$t = 0 ?$

Heaviside Step
func


~~u~~ $u(0) = \frac{1}{2}$

One of the most useful functions we use is the *unit step function*

$$u(t) = \begin{cases} 0 & t < 0 \\ 1 & t > 0 \end{cases} \quad (1)$$



Manipulating the unit step




We can do all the same things with step functions as with other functions:

- time shifts
- time reversal
- expansion/contraction



Interpreting/using the step function




- Making a signal “start” at a certain time:

$$f(t)u(t - \tau) = \begin{cases} f(t) & t > \tau \\ 0 & t < \tau \end{cases} \quad (2)$$

- “Switching on the system”: e.g. a DC voltage starts a time 1.
- Effect of integrating an impulse function (more on that later).




Example: piecewise constant functions



We can use step functions to build piecewise constant functions.



Square waves




Another example is a square wave: we make one pulse and then repeat it:

$$x_1(t) = u(t) - 2u(t-1) + u(t-2) \quad (3)$$

$$x(t) = \sum_{n=-\infty}^{\infty} x_1(t - 2n). \quad (4)$$



Rectangle function



A related function is the rectangle function:

$$\text{rect}(t) = \begin{cases} 1 & |t| < \frac{1}{2} \\ 0 & |t| > \frac{1}{2} \end{cases} \quad (5)$$

$$= u(t + 1/2) - u(t - 1/2). \quad (6)$$

Since it's so useful it gets its own notation.



Windowing functions


Rectangles are often used to *window* functions – this means only keeping the function inside a window. The function $\text{rect}(at - b)$ keeps a window of total length $\frac{1}{a}$ centered at $\frac{b}{a}$. So

$$f(t) \text{rect}(3t - 2) = \begin{cases} f(t) & \frac{1}{2} < t < \frac{5}{6} \\ 0 & \text{elsewhere} \end{cases} \quad (7)$$

$$\frac{b}{a} = \frac{2}{3} = \frac{4}{6}$$

$$\text{total } \frac{1}{3}$$

$\frac{1}{6}$ in each direction



Non-rectangular windows

In some applications a rectangular window is not desirable due to the sharpness of the cutoffs. You will see in MATLAB documentation references to all sorts of window functions. The rect function is called a “Hamming window.” Others include


- Hanning
- Kaiser
- Bartlett
- Parzen

There are a ton in the MATLAB documentation:

`https://www.mathworks.com/help/signal/windows.html`



The ramp function




If we integrate the unit step function we get the ramp function:

$$r(t) = \begin{cases} 0 & t \leq 0 \\ t & t > 0 \end{cases} \quad (8)$$

If you feed a step function into an integrator you'll get a ramp – this shows that the integrator is *unstable*.



Sawtooth waves




We can synthesize sawtooth waves with ramps and unit steps:

$$x_1(t) = 5r(t) - 5r(t - 2) - 10u(t - 2) \quad 0 \leq t \leq 2 \quad (9)$$

$$x(t) = \sum_{n=-\infty}^{\infty} x_1(t - 2n). \quad (10)$$



Wave synthesis: try it yourself




Problem

Write a formula for a triangular wave with period 6, minimum value 0 and maximum value 10. Try to plot it in MATLAB.



What about DT?




In DT we also use the unit step function

$$u[n] = \begin{cases} 1 & n \geq 0 \\ 0 & n < 0 \end{cases}$$

We don't tend to use the ramp and rect functions however.



What about DT?



We can use unit steps to make windowed versions of signals.



Recap

- introduced the unit step, unit ramp, and rect functions in CT
- introduced the DT unit step
- showed how to use steps to make signals start and stop at different times
- constructed infinite waves using steps and ramps

