

Linear Systems and Signals

Complex exponentials in CT

Anand D. Sarwate

Department of Electrical and Computer Engineering
Rutgers, The State University of New Jersey

2020



Learning objectives

The learning objectives for this section are:

- Write the general form of complex exponential functions in CT.
- Use Euler's formula to express sinusoids in terms of complex exponentials.
- Sketch special cases of real exponentials, exponentially modulated sinusoids, and real sinusoids.
- Calculate fundamental period and frequency for a complex periodic exponential
- Explain similarities and differences between CT complex exponentials.



Basic definition


A CT *complex exponential signal* is of the form

$$x(t) = Ce^{at} \quad (1)$$

where C and a are complex numbers. All the other things we will see will be special cases of this form.




Real signals



If $x(t) = Ce^{at}$ is a *real* signal, then C and a are real numbers. These should be pretty familiar looking: if $a > 0$ then there is an exponential increase and if $a < 0$ there is an exponential decrease.



Real signals



If $x(t) = Ce^{at}$ is a *real* signal, then C and a are real numbers. These should be pretty familiar looking: if $a > 0$ then there is an exponential increase and if $a < 0$ there is an exponential decrease.




Complex signals

If we make a purely imaginary then we have

$$x(t) = Ce^{j\omega_0 t} = |C|e^{j(\omega_0 t + \angle C)} \quad (2)$$

Let's focus on the simple case where $C = 1$. Then we have

$$x(t) = e^{j\omega_0 t}. \quad (3)$$



This is *periodic* with fundamental period

$$T_0 = \frac{2\pi}{|\omega_0|}. \quad (4)$$



Eulerizing and sinusoids

Euler's formula is going to be used *a lot* in this course:

$$e^{j\omega_0 t} = \cos(\omega_0 t) + j \sin(\omega_0 t) \quad (5)$$

What this implies is that regular sines and cosines are linear combinations of complex exponentials:

$$\cos(\omega_0 t) = \frac{e^{j\omega_0 t} + e^{-j\omega_0 t}}{2} \quad (6)$$


$$\sin(\omega_0 t) = \frac{e^{j\omega_0 t} - e^{-j\omega_0 t}}{2j} \quad (7)$$

$$\frac{1}{j} = -j$$

It's often very useful to “Eulerize” sines and cosines so that you can “gather like terms” when doing algebra. As an exercise, try getting a formula for $\cos(\omega_0 t) \sin(\omega_0 t)$ by Eulerizing, multiplying, and then simplifying back to real signals.



Now with phases!



If we have a phase shift:

$$A \cos(\omega_0 t + \phi) = \frac{A}{2} e^{j\phi} e^{j\omega_0 t} + \frac{A}{2} e^{-j\phi} e^{-j\omega_0 t} = A \Re \left\{ e^{j\omega_0 t + \phi} \right\} \quad (8)$$

Note that the amplitudes of each term are **complex-valued** (in our general formula, C is complex). Likewise:

$$A \sin(\omega_0 t + \phi) = A \Im \left\{ e^{j\omega_0 t + \phi} \right\}. \quad (9)$$



Harmonically related sinusoids

If we have a complex exponential $\phi_1(t) = e^{j\omega_0 t}$ with period $T_0 = \frac{2\pi}{|\omega_0|}$ we can define the *harmonically related* complex exponentials, which are

$$\phi_k(t) = e^{jk\omega_0 t} \quad k = 0, \pm 1, \pm 2, \dots \quad (10)$$

These are all of the periodic exponentials whose frequency is an integer multiple of $\phi_1(t)$.

This terminology comes from music (and the physics thereof): a vibrating string or standing wave is the superposition (linear combination) of harmonically related exponentials.



The general case

More generally, we can split C into its magnitude and phase form and a into its Cartesian form:

$$C = |C|e^{j\theta} \quad a = r + j\omega_0 \quad (11)$$

and look at functions of the form


$$x(t) = Ce^{at} = |C|e^{j\theta} e^{(r+j\omega_0)t} = |C|e^{rt} \underline{e^{j(\omega_0 t + \theta)}} \quad (12)$$

Eulerizing:

$$x(t) = |C|e^{rt} \underline{\cos(\omega_0 t + \theta)} + j|C|e^{rt} \underline{\sin(\omega_0 t + \theta)} \quad (13)$$



The general case in pictures



The general case in pictures

