

Linear Systems and Signals

Complex exponentials in DT

Anand D. Sarwate

Department of Electrical and Computer Engineering
Rutgers, The State University of New Jersey

2020



Learning objectives

The learning objectives for this section are:

- Write the general form of complex exponential functions in DT.
- Use Euler's formula to express sinusoids in terms of complex exponentials.
- Sketch special cases of real exponentials, exponentially modulated sinusoids, and real sinusoids.
- Calculate fundamental period and frequency for a complex periodic exponential
- Explain similarities and differences between DT complex exponentials.



The DT complex exponential sequence

We have a *similar* story in DT. A complex exponential signal is of the form


$$x[n] = C\alpha^n \quad C, \alpha \in \mathbb{C} \quad (1)$$

Why don't we write $\alpha = e^\beta$ where β is complex? Because it's often more convenient to write things in this form instead. That's because we will be doing a lot of geometric series:

$$\sum_{n=0}^{\infty} x[n] = C \sum_{n=0}^{\infty} \alpha^n = \frac{1}{1-\alpha} \quad |\alpha| < 1 \quad (2)$$



Real signals



Real DT exponentials can be monotonically increasing, monotonically decreasing, or alternating:

$$x[n] = (1.1)^n \quad (3)$$


$$(4)$$

$$(5)$$

$$(6)$$



Real signals



Real DT exponentials can be monotonically increasing, monotonically decreasing, or alternating:

$$x[n] = (1.1)^n \quad (3)$$


$$x[n] = (0.9)^n \quad (4)$$

$$(5)$$

$$(6)$$



Real signals



Real DT exponentials can be monotonically increasing, monotonically decreasing, or alternating:

$$x[n] = (1.1)^n \quad (3)$$


$$x[n] = (0.9)^n \quad (4)$$

$$x[n] = (-1.1)^n \quad (5)$$

$$(6)$$



Real signals



Real DT exponentials can be monotonically increasing, monotonically decreasing, or alternating:

$$x[n] = (1.1)^n \quad (3)$$


$$x[n] = (0.9)^n \quad (4)$$

$$x[n] = (-1.1)^n \quad (5)$$

$$x[n] = (-0.9)^n \quad (6)$$



General complex exponentials



We can write $C = |C|e^{j\theta}$ and $\alpha = |\alpha|e^{j\omega_0}$ in magnitude-phase form to get


$$x[n] = |C||\alpha|^n e^{j(\omega_0 n + \theta)} \quad (7)$$

$$= |C||\alpha|^n \cos(\omega_0 n + \theta) + j|C||\alpha|^n \sin(\omega_0 n + \theta) \quad (8)$$

The real/imaginary part look similar to the CT version.



General complex exponentials



We can write $C = |C|e^{j\theta}$ and $\alpha = |\alpha|e^{j\omega_0}$ in magnitude-phase form to get

$$x[n] = |C||\alpha|^n e^{j(\omega_0 n + \theta)} \quad (7)$$

$$= |C||\alpha|^n \cos(\omega_0 n + \theta) + j|C||\alpha|^n \sin(\omega_0 n + \theta) \quad (8)$$

The real/imaginary part look similar to the CT version.



Periodicity

Focusing on

$$x[n] = e^{j\omega_0 n} \quad (9)$$

we see that if we add/subtract any integer multiple of 2π to ω_0 we get the same thing:

$$e^{j(\omega_0 + 2\pi k)n} = e^{j\omega_0 n} e^{j2\pi k n} = e^{j\omega_0 n} 1^k = e^{j\omega_0 n}. \quad (10)$$

That means we can think of ω_0 as taking values in an interval of length 2π , such as $[0, 2\pi]$ or $[-\pi, \pi]$. For $x[n]$ to be *periodic* with period N we need

$$\omega_0 N = 2\pi m \quad \implies \quad \frac{\omega_0}{2\pi} = \frac{m}{N}. \quad (11)$$

← rational number



Periodicity

Not all DT complex exponentials are periodic! We have to have $\frac{2\pi}{\omega_0}$ be a rational number. The fundamental period of a complex exponential $e^{j\omega_0 n}$ is the smallest N such that there is an integer m with

$$N = m \left(\frac{2\pi}{\omega_0} \right). \quad (12)$$

Note, this is very different from CT, where $T_0 = \frac{2\pi}{|\omega_0|}$.

Example: (*OW p.29*) Suppose $x[n] = \cos(8\pi n/31)$ and $x(t) = \cos(8\pi t/31)$. Then

- $x(t)$ is periodic with period $\frac{2\pi}{8\pi/31} = \frac{31}{4}$.
- $x[n]$ is periodic with period 31.



Harmonically related exponentials

What about *harmonically related* exponentials in DT? The periodic complex exponential with period N has $\omega_0 = \frac{2\pi}{N}$. So

$$\phi_1[n] = e^{j(2\pi/N)n} \quad e^{j2\pi n} = 1 \quad (13)$$

The elements of $\phi_1[n]$ are also called the N -th roots of unity. We can define the *harmonically related* exponentials as those with integer multiples of ω_0 :

$$\phi_k[n] = e^{jk(2\pi/N)n} \quad (14)$$

Note: when k hits N we get $\phi_N[n] = e^{j2\pi n} = 1$ so $\phi_{N+1}[n] = \phi_1[n]$. If we define $\phi_0[n] = 1$ then there are only N unique harmonically related exponentials with period N .



An example

Take $N = 6$, so $\omega_0 = 2\pi/6 = \pi/3$:



n	0	1	2	3	4	5
$\phi_0[n]$	1	1	1	1	1	1
$\phi_1[n]$	1	$e^{j(\pi/3)n}$	$e^{j(2\pi/3)n}$	-1	$e^{j(4\pi/3)n}$	$e^{j(5\pi/3)n}$ ✓
$\phi_2[n]$	1	$e^{j(2\pi/3)n}$	$e^{j(4\pi/3)n}$	1	$e^{j(2\pi/3)n}$	$e^{j(4\pi/3)n}$
$\phi_3[n]$	1	-1	1	-1	1	-1
$\phi_4[n]$	1	$e^{j(4\pi/3)n}$	$e^{j(2\pi/3)n}$	1	$e^{j(4\pi/3)n}$	$e^{j(2\pi/3)n}$
$\phi_5[n]$	1	$e^{j(5\pi/3)n}$	$e^{j(4\pi/3)n}$	-1	$e^{j(2\pi/3)n}$	$e^{j(\pi/3)n}$

Here we make a few simplifications:

$$e^{j(3\pi/3)n} = -1 \quad (15)$$

$$e^{j(6\pi/3)n} = 1 \quad (16)$$

$$e^{j((m+6)\pi/3)n} = e^{j(m\pi/3)n} \quad (17)$$



Summary of differences between DT and CT

	$e^{j\omega_0 t}$	$e^{j\omega_0 n}$
range of ω_0	$\omega_0 \in \mathbb{R}$	$\omega_0 \in [0, 2\pi]$
is periodic?	for any ω_0	for $\omega_0 = \frac{2\pi m}{N}$
fundamental frequency	ω_0	ω_0/m
fundamental period	$T_0 = \frac{2\pi}{ \omega_0 }$	$\underline{N} = m \frac{2\pi}{\omega_0}$

