

Linear Systems and Signals

Computing the convolution using superposition

Anand D. Sarwate

Department of Electrical and Computer Engineering
Rutgers, The State University of New Jersey

2020



Learning objectives

The learning objective for this section is:

- calculate simple convolutions by superimposing outputs from impulses



Impulse response and convolution

$$y[n] = \sum_{k=-\infty}^{\infty} x[k] \underline{h[n-k]} = \sum_{k=-\infty}^{\infty} \underline{h[k]} x[n-k] \quad (1)$$

For “simple” signals and filters with “simple” impulse responses we can sometimes just compute the convolution directly:

- 1 Interpret the output as scaled and shifted copies of the impulse response $h[n]$ or as scaled and shifted copies of $x[n]$.
- 2 Use the “flip-and-slide” view to compute the product $h[k]x[n-k]$ for each n and add it up to get $y[n]$.
- 3 Write out the convolution formula and use formulas such as power series to simplify the expression for the output.



Direct calculation by superposition

Look at the formula

$$y[n] = \sum_{k=-\infty}^{\infty} x[k] h[n-k] \quad (2)$$

Handwritten notes:
 - "coefficients" with an arrow pointing to $x[k]$
 - " $h[n]$ delayed by k " with an arrow pointing to $h[n-k]$

This is a sum of scaled and shifted copies of $h[n]$. Sometimes we can just write these out. For example, take $x[n] = \delta[n] - \delta[n-1]$ and

$$h[n] = 3\delta[n] + 2\delta[n-1] - \delta[n-2]:$$

$$y[n] = 3\delta[n] + 2\delta[n-1] - \delta[n-2] \quad (3)$$

Handwritten notes:
 - "make one copy @ time $n=0$ " with an arrow pointing to $3\delta[n]$
 - "subtract one copy @ $n=1$ " with an arrow pointing to $-\delta[n-2]$

$$- 3\delta[n-1] - 2\delta[n-2] + \delta[n-3] \quad (4)$$

$$= 3\delta[n] - \delta[n-1] - 3\delta[n-2] + \delta[n-3] \quad (5)$$





Example

Find $y[n] = (x * h)[n]$ when

$$x[n] = u[n] - u[n - 4]$$

$$h[n] = u[n] - u[n - 3]$$

Step 0: draw a picture and rewrite the signals if needed:



$$x[n] = \begin{cases} 1 & 0 \leq n \leq 3 \\ 0 & \text{otherwise} \end{cases}$$

$$h[n] = \begin{cases} 1 & 0 \leq n \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

Direct algebraic approach

- ① Write $x[n]$ and $h[n]$ in terms of δ -functions:

$$x[n] = \delta[n] + \delta[n-1] + \delta[n-2] + \delta[n-3]$$

$$h[n] = \delta[n] + \delta[n-1] + \delta[n-2]$$



- ② Write scales and shifts of $h[n]$:

$$\begin{array}{ccccccc}
 y[n] = & \delta[n] + \delta[n-1] & + \delta[n-2] & & & & \\
 & + \delta[n-1] & + \delta[n-2] + \delta[n-3] & & & & \\
 & & + \delta[n-2] + \delta[n-3] & + \delta[n-4] & & & \\
 & & & + \delta[n-3] & + \delta[n-4] + \delta[n-5] & & \\
 & & & & & &
 \end{array}$$

Handwritten annotations: Blue arrows point from the terms to their corresponding shifts of $h[n]$. The shifts are labeled as $h[n]$, $h[n-1]$, $h[n-2]$, and $h[n-3]$.


- ③ Add them up:

$$\begin{aligned}
 y[n] = & \delta[n] + 2\delta[n-1] + 3\delta[n-2] + 3\delta[n-3] \\
 & + 2\delta[n-4] + \delta[n-5]
 \end{aligned}$$



Try some yourself

Problem



Find the convolution ~~from the following input output relations:~~

$$h[n] = u[n] - u[n - 4], x[n] = -\delta[n] + 2\delta[n - 2]$$

$$h[n] = n(u[n] - u[n - 3]), x[n] = 2\delta[n] - 2\delta[n - 1]$$

$$h[n] = \text{see figure above}$$

Make up a few on your own!