

Linear Systems and Signals

System properties, the ROC, and inverse systems

Anand D. Sarwate

Department of Electrical and Computer Engineering
Rutgers, The State University of New Jersey

2020



Learning objectives

The learning objectives for this section are:

- characterize stability and causality of a system in terms of the ROC
- find all time domain signals with a given transform
- determine if causal, stable inverses exist



System properties

We can relate system properties to the ROC of a rational transform $H(s)$ with ROC \mathcal{R}_h :

don't allow delay $e^{-s\tau}$

- If $H(s)$ is causal system then \mathcal{R}_h goes from the rightmost pole to ∞ .
 $h(t)$ is right-sided

- If $H(s)$ is anticausal system then \mathcal{R}_h goes from $-\infty$ to the leftmost pole.

- $H(s)$ is stable if and only if \mathcal{R}_h contains the imaginary axis $\{s = j\omega\}$.

- $H(s)$ is invertible if there is a $G(s)$ with ROC \mathcal{R}_g such that $H(s)G(s) = 1$ and $\mathcal{R}_h \cap \mathcal{R}_g \neq \emptyset$

$\frac{1}{H(s)}$


causal + stable

\Rightarrow all poles are in the left half-plane.

We can use these properties to find a time domain signal with a given Laplace transform.



Different signals, same transform, different ROCs



Recall that multiple signals can have the same Laplace transform with different ROCs. For example:

$$Y(s) = \frac{(s-1)}{(s+3)(s+1)} = \frac{2}{s+3} - \frac{1}{s+1} \quad (1)$$

Handwritten notes: zero @ 1, poles @ -3, -1

can come from

$$\text{right} \quad y_1(t) = 2e^{-3t}u(t) - e^{-t}u(t) \quad (2)$$

$$\text{left} \quad y_2(t) = -2e^{-3t}u(-t) + e^{-t}u(-t) \quad (3)$$

$$\text{two} \quad y_3(t) = 2e^{-3t}u(t) + e^{-t}u(-t) \quad (4)$$



Convolution and the ROCs

The convolution property says that if $h(t) \xleftrightarrow{\mathcal{L}} H(s)$ and $x(t) \xleftrightarrow{\mathcal{L}} X(s)$ then

$$y(t) = (h * x)(t) \xleftrightarrow{\mathcal{L}} Y(s) = H(s)X(s) \quad (5)$$

where the ROC contains $\mathcal{R}_x \cap \mathcal{R}_h$. Suppose we have

$$X(s) = \frac{s-1}{s+1} = 1 - \frac{2}{s+1} \quad (6)$$

$$H(s) = \frac{1}{s+3} \quad (7)$$

$$Y(s) = \frac{(s-1)}{(s+3)(s+1)} \quad (8)$$



Four options?

We have two different $x(t)$ signals with the same $X(s)$ but different \mathcal{R}_x . Likewise, we have two different $h(t)$ signals with the same $H(s)$.

	$x(t)$	<i>right</i> $\delta(t) - 2e^{-t}u(t)$	<i>left</i> $\delta(t) + 2e^{-t}u(-t)$
$h(t)$			
<i>right</i> $e^{-3t}u(t)$		right-sided $y_1(t)$	two-sided $y_3(t)$
<i>left</i> $-e^{-3t}u(-t)$		<u>no intersection</u>	<u>left-sided $y_2(t)$</u>

In order for $y(t)$ to exist the intersection $\mathcal{R}_x \cap \mathcal{R}_h$ cannot be empty.



Example: finding the ROCs

Let's find all the impulse responses which can lead to a transform

$$H(s) = \frac{s + 5}{(s + 2)(s + 4)(s + 6)} \quad (9)$$

This has a zero at $s = \underline{-5}$ and poles at $s = \underline{-2}$, $s = \underline{-6}$, and $s = \underline{-4}$.

Possible ROCs:

$$\checkmark \{ \Re\{s\} < -6 \} \quad \text{left-sided, unstable} \quad (10)$$


$$\checkmark \{ -6 < \Re\{s\} < -4 \} \quad \text{two-sided, unstable} \quad (11)$$

$$\checkmark \{ -4 < \Re\{s\} < -2 \} \quad \text{two-sided, unstable} \quad (12)$$

$$\checkmark \{ \Re\{s\} > \underline{-2} \} \quad \text{right-sided, stable} \quad (13)$$



The pole-zero diagram



Split using partial fraction expansion

Using partial fraction expansion:

$$\frac{s+5}{(s+4)(s+6)} \Big|_{s=-2} = \frac{3}{8} \quad \checkmark \quad (14)$$

$$\frac{s+5}{(s+2)(s+6)} \Big|_{s=-4} = -\frac{1}{4} \quad \checkmark \quad (15)$$

$$\frac{s+5}{(s+2)(s+4)} \Big|_{s=-6} = -\frac{1}{8} \quad \checkmark \quad (16)$$

So


$$H(s) = \frac{3/8}{s+2} - \frac{1/4}{s+4} - \frac{1/8}{s+6} \quad (17)$$

$\xrightarrow{A_1}$ $\xrightarrow{A_2}$ $\xrightarrow{A_3}$

So for each of the corresponding ROCs we have to choose which inverse (left- or right-sided) time-domain signal to choose.



Associating the time-domain signals



$$\begin{aligned} &\leftarrow e^{-2t} u(-t) \\ &\leftarrow e^{-4t} u(-t) \\ &e^{-6t} u(t) \rightarrow \end{aligned}$$

$$h(t) = -\frac{3}{8}e^{-2t}u(-t) + \frac{1}{4}e^{-4t}u(-t) + \frac{1}{8}e^{-6t}u(-t)$$

left-sided

$$\{\Re\{s\} < -6\}$$

$$h(t) = -\frac{3}{8}e^{-2t}u(-t) + \frac{1}{4}e^{-4t}u(-t) - \frac{1}{8}e^{-6t}u(t)$$

$$\{-6 < \Re\{s\} < -4\}$$

$$h(t) = -\frac{3}{8}e^{-2t}u(-t) - \frac{1}{4}e^{-4t}u(t) - \frac{1}{8}e^{-6t}u(t)$$

$$\{-4 < \Re\{s\} < -2\}$$

$$h(t) = \frac{3}{8}e^{-2t}u(t) - \frac{1}{4}e^{-4t}u(t) - \frac{1}{8}e^{-6t}u(t)$$

right-sided

$$\{\Re\{s\} > -2\}$$



Determining inverses

Does the system with the following impulse response have a stable, causal inverse?

$$H(s) = \frac{(s+1)(s+2)}{s+3} \quad \mathcal{R}_h = \{\Re\{s\} > -3\} \quad (18)$$

Let's take the algebraic inverse:

$$\frac{1}{H(s)} = G(s) = \frac{(s+3)}{(s+1)(s+2)} = \frac{2}{s+1} - \frac{1}{s+2} \quad (19)$$

This has 3 ROCs: $\{\Re\{s\} < -2\}$, $\{-2 < \Re\{s\} < -1\}$, and $\{\Re\{s\} > -1\}$. The last one is stable and causal, and

contains imag axis \Rightarrow stable

$$\{\Re\{s\} > -3\} \cap \{\Re\{s\} > -1\} = \{\Re\{s\} > -1\} \quad (20)$$

So $g(t) = 2e^{-t}u(t) - e^{-2t}u(t)$ is a stable causal inverse.



Sometimes inverses are not realizable

Does the system with the following impulse response have a stable, and causal inverse?

$$H(s) = \frac{(s - 3)}{(s + 1)(s + 2)}$$

$$\mathcal{R}_h = \{\Re\{s\} > -1\} \quad (21)$$

Taking the algebraic inverse:

$$G(s) \stackrel{= 1/H(s)}{=} \frac{(s + 1)(s + 2)}{(s - 3)} = \frac{s^2 + 3s + 2}{s - 3} \stackrel{\text{long division}}{=} s + 6 - \frac{16}{s - 3} \quad (22)$$

This has 2 ROCs: $\{\Re\{s\} < 3\}$, $\{\Re\{s\} > 3\}$. We have two choices:

$$\{-1 < \Re\{s\} < 3\}$$

$$\{\Re\{s\} > 3\}$$

stable, two-sided

unstable, right-sided

So the answer is no.



Try it yourself

Problem

Find all time domain signals with the following Laplace transforms and determine if the systems are causal/anticausal or stable/unstable:

$$H(s) = \frac{s - 3}{s^2 + 5s + 6} \quad (23)$$

$$H(s) = \frac{s - 1}{2s^2 + 11s + 12} \quad (24)$$

$$H(s) = \frac{s + 6}{s^2 + 2s - 15} \quad (25)$$

$$H(s) = \frac{s + 2}{s^3 + 3s^2 - s - 3} \quad (26)$$

Determine if there are causal, stable inverses for each of them.

